

International Conference on Magnetism and
Superconductivity in Topological Matter

NOV 9-11, 2023 HYBRID MEETING

REGISTER TODAY!

Science
MAAS



SAS
北京国际学术交流会

中关村论坛
2023 SGC FORUM

Pavlovian conditioning–induced hallucinations result from over- weighting of perceptual priors

A. R. POWERS¹, C. MATHYS¹, AND P. R. CORLETT¹ [Authors Info & Affiliations](#)

SCIENCE • 11 Aug 2017 • Vol 357, Issue 6351 • pp. 596-600

5,467



Neural mechanisms for hallucinations

Pairing a stimulus in one modality (vision) with a stimulus in another (sound) can lead to task-induced hallucinations in healthy individuals. After many trials, people eventually report perceiving a nonexistent stimulus contingent on the presence of the previously paired stimulus. Powers *et al.* investigated how different groups of volunteers and patients respond to this conditioning paradigm. They used behavior, neuroimaging, and computational modeling to dissect the effect of perceptual priors versus sensory evidence on such induced hallucinations. People who are more prone to hear voices were more susceptible to the induced auditory hallucinations. The network of brain regions that was active during the conditioned hallucinations resembled the network observed during clinical symptom capture in individuals who hallucinate while in a brain scanner.

Science, this issue p. [596](#)

Abstract

Some people hear voices that others do not, but only some of those people seek treatment. Using a Pavlovian learning task, we induced conditioned hallucinations in four groups of people who differed orthogonally in their voice-hearing and treatment-seeking statuses. People who hear voices were significantly more susceptible to the effect. Using functional neuroimaging and computational modeling of perception, we identified processes that differentiated voice-hearers from non-voice-hearers and treatment-seekers from non-treatment-seekers and characterized a brain circuit that mediated the conditioned hallucinations. These data demonstrate the profound and sometimes pathological impact of top-down cognitive processes on perception and may represent an objective means to discern people with a need for treatment from those without.

Perception is not simply the passive reception of inputs (1). We actively infer the causes of our sensations (2). These inferences are influenced by our prior experiences (3). Priors and inputs might be combined according to Bayes' rule (4). Prediction errors, the mismatch between priors and inputs, contribute to belief updating (5). Hallucinations (percepts without external stimulus) may arise when strong priors cause a percept in the absence of input (6). We tested this theory by engendering new priors about auditory stimuli in human observers using Pavlovian conditioning.

Even in healthy individuals, the repeated co-occurrence of visual and auditory stimuli can induce auditory hallucinations (7). We examined this effect with functional imaging. Some argue that in patients with psychosis, weak priors lead to aberrant prediction errors, resulting in auditory verbal hallucinations (AVH) (8). Others have observed strong priors in patients, but the effects were not specific to hallucinations (9, 10). Such inconsistencies may reflect the hierarchical organization of perception: Perturbations may affect some levels of the hierarchy and not others (9). We used computational modeling to infer the strength of participants' hierarchical perceptual beliefs from their behavioral responses during conditioning (11). Our model captured how priors are combined with sensory evidence, allowing us to test the strong-prior hypothesis directly.

Participants worked to detect a 1-kHz tone occurring concurrently with presentation of a checkerboard visual stimulus. First, we determined individual thresholds for detection and psychometric curves (12). Then, at the start of conditioning, the tone was presented frequently at threshold (Fig. 1A, left), engendering a belief in audio-visual association. This belief was then tested (Fig. 1A, right) with increasingly frequent subthreshold and target-absent trials (Fig. 1B). Conditioned hallucinations occurred when subjects reported tones that were not presented, conditional upon the visual stimulus.




Fig. 1 Methods and behavioral results.

(A) Trials consisted of simultaneous presentation of a 1000-Hz tone in white noise and a visual checkerboard. (B) We estimated individual psychometric curves for tone detection (left) and then systematically varied stimulus intensity over 12 blocks of 30 conditioning trials. Threshold tones were more likely early, and absent tones were more likely later (right). (C) Groups varied along two dimensions: the presence (+) or absence (-) of daily AVH (blue) and the presence (+) or absence (-) of a diagnosable psychotic-spectrum illness (red). (D) Detection thresholds. Error bars represent ± 1 SD, and boxes represent ± 1 SEM. (E) Probability of conditioned hallucinations varied according to hallucination status. Error bars represent ± 1 SD, and boxes represent ± 1 SEM. (Inset) Error bars represent ± 1 SEM. $***P < 0.001$. (F) Differences between hallucinating and nonhallucinating groups were found only in the target-absent and 25% likelihood of detection conditions. Error bars represent ± 1 SEM. (G) Hallucinators were more confident than nonhallucinators when reporting a tone that did not exist. $*P < 0.05$. (H and I) Both the probability of reporting conditioned hallucinations (H) and the confidence with which they were reported (I) correlated with a measure of hallucination severity.

We recruited four groups of subjects (Fig. 1C): people with a diagnosed psychotic illness who heard voices (P+H+, $n = 15$); those with a similar illness who did not hear voices (P+H-, $n = 14$); an active control group who heard daily voices, but had no diagnosed illness (P-H+, $n = 15$) (13)—they attributed their experiences metaphysically (supplementary materials) (14); and last, controls without diagnosis or voices (P-H-, $n = 15$).

Groups were matched demographically (tables S1 to S4). Rates of detection of tones at threshold were similar across groups. All groups demonstrated conditioned hallucinations. However, those with daily hallucinations endorsed more conditioned hallucinations than those without, regardless of diagnosis ($F_{1,55} = 19.59$, $P = 5.82 \times 10^{-5}$) (Fig. 1D). This effect remained after accounting for differences in detection thresholds (Fig. 1E, fig. S1, and table S5). Group differences in propensity to report tones were observed only in the “no-tone” and 25% “likelihood of detection” conditions (intensity-by-hallucination status $F_{3,165} = 13.59$, $P = 5.73 \times 10^{-4}$) (Fig. 1F).

Participants also rated their decision confidence by holding down the response button (Fig. 1G). Participant confidence varied with stimulus intensity (“yes”: $R = 0.39$, $P = 7.46 \times 10^{-10}$; “no”: $R = 0.22$, $P = 9.02 \times 10^{-4}$). However, hallucinators were more confident in their conditioned hallucinations than non-hallucinators ($F_{1,53} = 6.50$, $P = 0.045$). Both conditioned hallucinations and confidence correlated with

hallucination severity outside of the laboratory ([Fig. 1, H and I](#), and fig. S3).

In order to establish whether conditioned hallucinations involved true percepts, we first identified tone-responsive regions from thresholding runs [peaks at (-60, -20, 2) and (62, -28, 10)] ([Fig. 2A](#)). As observed with elementary hallucinations ([15](#)), activity in tone-responsive regions was greater during conditioned hallucinations compared with correct rejections ($t_{56} = 4.93, P = 7.59 \times 10^{-6}$) ([Fig. 2B](#)). Electrical stimulation of this region in human patients produces AVH ([16](#)). Taken together, these findings are consistent conditioned hallucinations involving actual perception.




Fig. 2 Imaging results.

(A) Bilateral supplemental auditory cortex covaried with tone intensity during thresholding (family-wise error rate-corrected, $P < 0.05$). (B) Parameter estimates from this region showed increased activation during conditioned hallucinations. $***P < 0.001$. (C) Whole-brain analysis during conditioned hallucinations (false discovery rate-corrected, $P < 0.05$). (D) Clusters derived from a meta-analysis ([17](#)) of AVH experiences during functional imaging. (E and F) Hallucinators were much less likely to engage ACC during correct rejections. Error bars represent ± 1 SEM.

Whole-brain analysis revealed that conditioned hallucinations also engaged anterior insula cortex (AIC), inferior frontal gyrus, head of caudate, anterior cingulate cortex (ACC), auditory cortex, and posterior superior temporal sulcus (STS) ([Fig. 2C](#) and table S6). A meta-analysis of symptom-capture-

based studies examining neural activity of AVH highlighted similar regions ([Fig. 2D](#)) ([17](#)). AIC and ACC responses frequently correlate with stimulus salience ([18](#)). However, their activation before near-threshold stimulus presentation predicts detection ([19](#)). Caudate is engaged during audiovisual associative learning ([20](#)). Likewise, AIC and ACC are engaged during multisensory integration ([21](#)).

There were no significant between-group differences in brain responses during conditioned hallucinations. However, hallucinators deactivated ACC more [peak at (-16, 54, 14); cluster-extent thresholded, starting value 0.005, critical cluster extent (k_e) = 99] during correct rejections compared with nonhallucinators ([Fig. 2, E and F](#)).

To further dissect conditioned hallucinations, we modeled their underlying computational mechanisms ([Fig. 3A](#)) using the hierarchical Gaussian filter (HGF) ([11](#)). We defined a perceptual model consisting of low-level perceptual beliefs (X_1), visual-auditory associations (X_2), and the volatility of those associations (X_3), as well as evolution rates encoding the relationships between levels (ω , θ). Critically, our perceptual model allowed for variability in weighting between sensory evidence and perceptual beliefs (v). For $v = 1$, prior and observation have equal weight; for $v > 1$, the prior has more weight than that of the observation (strong priors); and for $v < 1$, the observation has more weight than that of the prior (weak priors). The resultant posterior probability of a tone is then fed to a separate response model.




Fig. 3 HGF analysis.

(A) Computational model, mapping from experimental stimuli to observed responses through perceptual and response models. The first level (X_1) represents whether the subject believes a tone was present or not on trial t . The second level (X_2) is their belief that visual cues are associated with tones. The third level (X_3) is their belief about the volatility of the second level. The HGF allows for individual variability in weighting between sensory evidence and perceptual beliefs (parameter v). (B) At X_3 , there was a significant block-by-psychosis interaction. $*P < 0.05$. (C and D) Significant block-by-hallucination status interactions were seen at layers (D) X_1 and (C) X_2 . $***P < 0.001$. (E) v was significantly higher in those with hallucinations when compared with their nonhallucinating counterparts. $***P < 0.001$. (F) No main effects of group or interaction effects were seen for the decision noise parameter within the response model. Error bars and line shadings represent ± 1 SEM. Purple, P+H+; blue, P-H+; red, P+H-; white, P-H-.

Model parameters were fit to behavioral data, and the model was optimized by using log model evidence and simulations of observed behavior (figs. S3 and S4). Mean trajectories of perceptual beliefs were compared across groups (Fig. 3, B to D). Participants with hallucinations exhibited stronger beliefs at levels 1 (X_1 : $F_{11,605} = 4.8$, $P = 3.89 \times 10^{-7}$) (Fig. 3D) and 2 (X_2 : $F_{11,605} = 3.89$, $P = 1.84 \times 10^{-5}$) (Fig. 3C). X_3 beliefs evolved less in those with psychosis, who failed to recognize the increasing volatility in contingencies ($F_{11,605} = 2.11$, $P = 0.018$) (Fig. 3A).

Consistent with strong-prior theory, v was significantly larger in those with hallucinations when com-

pared with their nonhallucinating counterparts ([Fig. 3E](#)), regardless of diagnosis ($F_{1,55} = 13.96, P = 4.45 \times 10^{-4}$). Response model parameters did not differ across the groups ([Fig. 3F](#)).

We regressed model parameters onto task-induced brain responses ([Fig. 4A](#)). The X_1 trajectory covaried with several conditioned hallucination-responsive regions, including STS (table S7). X_3 trajectories, by contrast, covaried with hippocampus/parahippocampal gyrus and medial cerebellum (table S8). Parameter estimates from the X_1 -sensitive STS $[(-46 \ -36, 0), T_{57} = 2.09, P = 0.042]$ ([Fig. 4B](#)) and AIC $[(36, 8, -8), T_{57} = 2.26, P = 0.027]$ ([Fig. 4C](#)) were significantly greater in those with hallucinations versus those without. This is consistent with STS conferring auditory expectations that are responsive to incoming visual input ([22](#)). Parameter estimates from the X_3 -responsive cerebellar vermis $[(-2, -52, -16)]$ ([Fig. 4D](#)) were lower in participants with psychosis as compared with those without ($T_{57} = 2.05, P = 0.045$). In the model, subjects with psychosis were significantly less sensitive to the changes in contingency as the task progressed. Psychotic symptoms are often associated with pathological rigidity. Belief-updating correlated with responses in the hippocampus and cerebellum. Hippocampal activity correlates with uncertainty in perceptual predictions ([23](#)). The cerebellum has likewise been associated with production and updating of predictive models ([24](#)).




Fig. 4 HGF imaging results.

(A) HGF trajectories for X_1 (blue) and X_3 (red) regressed onto blood oxygen level-dependent time courses for the conditioned hallucinations task. Regions that identified significantly active during conditioned hallucinations (from Fig. 3C) are highlighted in yellow for reference. All images are cluster-extent thresholded at starting value 0.05; critical k_e for $X_1 = 545$ and $X_3 = 406$. (B and C) Parameter estimates of X_1 fit extracted from 5-mm sphere centered on (B) STS and (C) anterior insula activation differ based on hallucination status. (D) Parameter estimates of X_3 fit extracted from 1-mm sphere centered on cerebellar vermis activation differ based on psychosis status. Error bars represent 1 SEM.

Our X_1 , X_2 , and v findings are consistent with a strong-prior theory of hallucinations. The X_3 findings in psychotic patients may reflect a strong prior that contingencies are fixed. On the other hand, they could reflect a weak prior on volatility. These beliefs were not associated with hallucinations but rather psychosis more broadly. Under chronic uncertainty, secondary to consistent belief violation, it may be adaptive to resist updating beliefs (25).

Consistent with previous work applying signal detection theory (SDT) to AVH (26), we found liberal

criteria and low perceptual sensitivity in our H+ groups. A liberal criterion may reflect poor reality monitoring (26). However, meta-d' (a metric of participants' meta-cognitive sensitivity) did not differ significantly between groups (fig. S6). SDT is a descriptive tool that does not distinguish aberrant perceptions from decisions. Our modeling work, however, localized group differences to the perceptual model alone. The prior weighting parameter (ν) distinguished H+ from H- groups and also predicted confidence in conditioned hallucinations (fig. S7). Our observations support an explanation of hallucinations based on strong perceptual priors. They suggest precision treatments for hallucinations, such as targeting cholinergically mediated priors (27), and interventions to mollify psychosis more broadly, such as cerebellar transcranial magnetic stimulation (28).

Acknowledgments

The authors dedicate this work to the memory and legacy of Ralph E. Hoffman, M.D. Additional thanks go to M. Kelley, A. Bianchi, S. Bhatt, and E. Feeney for technical assistance as well as A. Nidiffer, L. Marks, S. Woods, and J. Krystal for their advice. This work was supported by the Connecticut Mental Health Center (CMHC) and Connecticut State Department of Mental Health and Addiction Services (DMHAS). P.R.C. was funded by an International Mental Health Research Organization/Janssen Rising Star Translational Research Award; National Institute of Mental Health (NIMH) grant 5R01MH067073-09; Clinical and Translational Science Award grant UL1 TR000142 from the National Center for Research Resources (NCRR) and the National Center for Advancing Translational Science (NCATS), components of the National Institutes of Health (NIH); NIH roadmap for Medical Research; the Clinical Neurosciences Division of the U.S. Department of Veterans Affairs; and the National Center for Post-Traumatic Stress Disorders, VA Connecticut Healthcare System (VACHS), West Haven, CT, USA. The contents of this work are solely the responsibility of the authors and do not necessarily represent the official view of NIH or the CMHC/DMHAS. A.R.P. was supported by the Integrated Mentored Patient-Oriented Research Training (IMPORT) in Psychiatry grant (5R25MH071584-07) as well as the Clinical Neuroscience Research Training in Psychiatry grant (5T32MH19961-14) from NIMH and a VA Schizophrenia Research Special Fellowship from VACHS, West Haven, CT, USA. Additional support was provided by the Yale Detre Fellowship for Translational Neuroscience as well as the Brain and Behavior Research Foundation in the form of a National Alliance for Research on Schizophrenia and Depression Young Investigator Award for A.R.P. The authors declare no conflicts of interest. Model code and data are stored at ModelDB (<http://modeldb.yale.edu/229278>). Imaging data are stored at NeuroVault (<http://neurovault.org/collections/OCFEJCQE/>).

Supplementary Material

Summary

Materials and Methods

Supplemental Text

Figs. S1 to S7

Tables S1 to S8

References (29, 30)

—

Resources

File (aan3458-powers-sm.pdf)

[DOWNLOAD](#)

1.16 MB

References and Notes

- 1 H. v. Helmholtz, *Treatise on Physiological Optics* (Voss, Hamburg, ed. 3, 1909).

[↶ GO TO REFERENCE](#) • [GOOGLE SCHOLAR](#)

- 2 K. Friston, A theory of cortical responses. *Philos. Trans. R. Soc. London B Biol. Sci.* **360**, 815–836 (2005).

[↶ GO TO REFERENCE](#) • [CROSSREF](#) • [PUBMED](#) • [GOOGLE SCHOLAR](#)

[SHOW ALL REFERENCES](#)

eLetters (1)

eLetters is a forum for ongoing peer review. eLetters are not edited, proofread, or indexed, but they are screened. eLetters should provide substantive and scholarly commentary on the article. Embedded figures cannot be submitted, and we discourage the use of figures within eLetters in general. If a figure is essential, please include a link to the figure within the text of the eLetter. Please read our [Terms of Service](#) before submitting an eLetter.

[LOG IN TO SUBMIT A RESPONSE](#)

DEC. 20, 2017

Hallucinations from overweighting of perceptual priors: Just the tip of the iceberg?

BARNABY NELSON Professor, Orygen, The National Centre of Excellence in Youth Mental Health, University of Melbourne

JESSICA ANNA HARTMANN Postdoctoral Research Fellow, Orygen, The National Centre of Excellence in Youth Mental Health, University of Melbourne

Powers et al's (1) important study demonstrates the role of top-down cognitive processes (i.e., implicit predictions) in shaping the incoming stream of sensory information. Their four-group design and use of experimentally-induced hallucinations convincingly shows that these aberrant predictive processes may play a role in t...

[view more](#)

Recommended articles from TrendMD

Cortical Map Reorganization Enabled by Nucleus Basalis Activity

Michael P. Kilgard et al., *Science*, 1998

Intonational speech prosody encoding in the human auditory cortex

C. Tang et al., *Science*, 2017

Renormalization Group Theory of Eigen Microstates [↗](#)

Teng Liu et al., *Chinese Physics Letters*, 2022

Variational quantum simulation of thermal statistical states on a superconducting quantum processor [↗](#)

Xue-Yi Guo et al., *Chinese Physics B*, 2023

Computational and neurobiological foundations of leadership decisions

Micah G. Edelson et al., *Science*, 2018

Restoring auditory cortex plasticity in adult mice by restricting thalamic adenosine signaling

Jay A. Blundon et al., *Science*, 2017

Synapse-specific representation of the identity of overlapping memory engrams

Kareem Abdou et al., *Science*, 2018

Bioinspired tactile perception platform with information encryption function [↗](#)

Zhi-Wen Shi et al., *Chinese Physics B*, 2022

Effects of Quantum Noise on Quantum Approximate Optimization Algorithm [↗](#)

Cheng Xue et al., *Chinese Physics Letters*, 2021

Total transmission from deep learning designs [↗](#)

Wu et al., *Journal of Electronic Science and Technology*, 2021

Powered by **TREND MD**




I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). [Learn more](#)

Yes

No

CURRENT ISSUE



Demographic and hormonal evidence for menopause in wild chimpanzees

BY BRIAN M. WOOD, JACOB D. NEGREY, ET AL.

Leveraging investment in agricultural science, technology, and innovation in Latin America and the Caribbean

BY EUGENIA SAINI

Universal scaling of the dynamic BKT transition in quenched 2D Bose gases

BY SHINICHI SUNAMI, VIJAY PAL SINGH, ET AL.

[TABLE OF CONTENTS >](#)

ADVERTISEMENT

Sign up for ScienceAdviser

Subscribe to *ScienceAdviser* to get the latest news, commentary, and research, free to your inbox daily.

[SUBSCRIBE >](#)

LATEST NEWS

NEWS | 2 NOV 2023

[Synthetic 'super melanin' speeds skin repair](#)

NEWS | 1 NOV 2023

[Heart-brain link offers new potential explanation for fainting](#)

NEWS | 1 NOV. 2023

[The universe's puzzlingly fast expansion may defy explanation, cosmologists fret](#)

NEWS | 1 NOV 2023

[Why don't fish have tonsils? They have a good alternative, study suggests](#)

NEWS | 1 NOV 2023

[Should scientists include their race, gender, or other personal details in papers?](#)

NEWS | 1 NOV 2023

[Where did shoulders come from? Ancient fish reveal joint's evolution](#)

ADVERTISEMENT

RELATED JOBS

Application Engineer Intern

Henkel
Singapore, Singapore

Assistant Professor of Biomedical Sciences

Cummings School of Veterinary Medicine at Tufts University, Department of Comparative Pathobiology
Grafton, Massachusetts

ADVERTISEMENT

Assistant or Associate Professor, Molecular Microbiology

Department of Microbiology, Ohio State University
Columbus, Ohio

[MORE JOBS](#) 

RECOMMENDED

REPORT | AUGUST 1973

Pavlovian Conditioning with Heat Reinforcement Produces Stimulus-Directed Pecking in Chicks

TECHNICAL COMMENT | JULY 1976

Conditioning of *Pleurobranchaea*

REPORT | FEBRUARY 1984

Epinephrine Enables Pavlovian Fear Conditioning Under Anesthesia

REPORT | DECEMBER 1965

Differential Classical Conditioning: Verbalization of Stimulus Contingencies

[View full text](#) | [Download PDF](#)

Science

Science
Advances

Science
Immunology

Science
Robotics

Science
Signalin

FOLLOW US



GET OUR NEWSLETTER

NEWS

[All News](#)
[ScienceInsider](#)
[News Features](#)
[Subscribe to News from Science](#)
[News from Science FAQ](#)
[About News from Science](#)

CAREERS

[Careers Articles](#)
[Find Jobs](#)
[Employer Hubs](#)

COMMENTARY

[Opinion](#)
[Analysis](#)
[Blogs](#)

JOURNALS

[Science](#)
[Science Advances](#)
[Science Immunology](#)
[Science Robotics](#)
[Science Signaling](#)
[Science Translational Medicine](#)
[Science Partner Journals](#)

AUTHORS & REVIEWERS

[Information for Authors](#)
[Information for Reviewers](#)

LIBRARIANS

[Manage Your Institutional Subscription](#)
[Library Admin Portal](#)
[Request a Quote](#)
[Librarian FAQs](#)

ADVERTISERS

[Advertising Kits](#)
[Custom Publishing Info](#)
[Post a Job](#)

RELATED SITES

[AAAS.org](#)
[AAAS Communities](#)
[EurekAlert!](#)
[Science in the Classroom](#)

ABOUT US

[Leadership](#)
[Work at AAAS](#)
[Prizes and Awards](#)

HELP

[FAQs](#)
[Access and Subscriptions](#)
[Order a Single Issue](#)
[Reprints and Permissions](#)
[TOC Alerts and RSS Feeds](#)
[Contact Us](#)



© 2023 American Association for the Advancement of Science. All rights reserved. AAAS is a partner of HINARI, AGORA, OARE, CHORUS, CLOCKSS, CrossRef and COUNTER.
Science ISSN [0036-8075](#)

[Terms of Service](#) | [Privacy Policy](#) | [Accessibility](#)